

Algebraic Number Theory

Problem sheet 8

- (3 points) Let K be a complete field with respect to an archimedean valuation $|\cdot|$. Show that K is isomorphic to either \mathbb{R} or \mathbb{C} such that the isomorphism induces a homeomorphism in the topology of K induced by $|\cdot|$.
- (3 points) Let K/\mathbb{Q} be a finite extension. Show that any nontrivial absolute value on K is equivalent to either the \mathfrak{p} -adic absolute value for some prime ideal $\mathfrak{p} \triangleleft \mathcal{O}_K$ (nonarchimedean case) or to the restriction of the usual absolute value of \mathbb{C} via an embedding $\tau: K \rightarrow \mathbb{C}$ (archimedean case). Two such absolute values are equivalent if and only if both are archimedean and the two corresponding embeddings $\tau_1, \tau_2: K \rightarrow \mathbb{C}$ are the complex conjugate of each other.
- (1 points) Write -1 as a p -adic integer $-1 = \sum_{i=0}^{\infty} a_i p^i$ ($a_i \in \{0, 1, \dots, p-1\}$).
- (2 points) Write $2/3$ and $-2/3$ in 5-adic form.
- (4 points) Verify that a p -adic number of the form $\sum_{i=-m}^{\infty} a_i p^i$ ($a_i = 0, 1, \dots, p-1, i \geq -m$) is rational if and only if the sequence of its digits is eventually periodic.
- (3 points) Solve the equation $x^2 = 2$ in \mathbb{Z}_7 .
- (3 points) Show that any automorphism of the field \mathbb{Q}_p is continuous, in particular equals the identity.
- (2 points) Let $n \geq 1$ be an integer and $n = a_0 + a_1 p + \dots + a_r p^r$ in base p ($0 \leq a_i < p$). Further put $s = a_0 + a_1 + \dots + a_r$. Verify that $v_p(n!) = \frac{n-s}{p-1}$.
- (2 points) Show that the sequence $1, 1/10, 1/100, \dots, 1/10^n, \dots$ does not converge in \mathbb{Q}_p for any prime p .
- (3 points) Let $\varepsilon \in 1 + p\mathbb{Z}_p$ and $\alpha = a_0 + a_1 p + a_2 p^2 + \dots$ be a p -adic integer with n th partial sum $s_n = a_0 + a_1 p + \dots + a_n p^n \in \mathbb{Z}$. Show that the limit $\varepsilon^\alpha := \lim_{n \rightarrow \infty} \varepsilon^{s_n}$ exists in \mathbb{Z}_p making $1 + p\mathbb{Z}_p$ a \mathbb{Z}_p -module (written multiplicatively).
- (3 points) Assume $(a, p) = 1$ ($a \in \mathbb{Z}$). Show that the sequence a^{p^n} converges in \mathbb{Q}_p .
- (2 points) Show that the fields \mathbb{Q}_p and \mathbb{Q}_q are not isomorphic for primes $p \neq q$.
- (2 points) Verify that the algebraic closure of \mathbb{Q}_p is an infinite extension of \mathbb{Q}_p .
- (5 points) Let $f(X) = a_0 + a_1 X + \dots + a_n X^n + \dots \in \mathbb{Z}_p[[X]]$ be a formal power series with coefficients in the ring of p -adic integers. Verify that f is convergent on the p -adic open unit disc $\{|X|_p < 1\}$ and has at most finitely many roots therein.